

Evaluation Of In-Vitro Anticancer Activity from Rhizophora Mucronata Mangrove Leaf Extracts Against Lung Cancer Cell Line

Aksha Sharen¹, Lakshmi T²

¹Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, (SIMATS), Savitha University, Chennai-77 India.

Email: 151801083.sdc@saveetha.com

²Professor, Department of Pharmacology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai-77 India.

Corresponding author
Lakshmi T

³Professor, Department of Pharmacology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai-77 India

Email: lakshmi@saveetha.com

Abstract

INTRODUCTION Herbal medicines play an important role in cancer prevention. Mangroves have been used worldwide due to the presence of various bioactive metabolites in it. The specific medicinal properties of Rhizophora mucronata have been an interest in preventing breast cancer. Mangroves Rhizophora mucronata belongs to the family Rhizophoraceae. The leaf extract reveals various roles in folk remedies to treat various diseases. The aim of the study is to determine anticancer activity of Rhizophora mucronata mangroves leaf extract against lung cancer cell line. **METHODOLOGY** The Rhizophora mucronata mangrove leaf samples were collected. The collected sample was washed and then dried for and turned into fine powder. A 25g dried powdered Rhizophora mucronata sample was mixed with 100ml of Ethanol and allowed to place for 24 hours at ambient temperature. Then the mixture was passed through whatman filter paper then the filtrate was centrifuged at 3000rpm for 10min and further filtered by 0.45µm syringe micro filter. At last, the solvents are evaporated via vacuum rotary evaporator until samples obtain in powder form. Then the sample was stored in a shadowy aluminum container at 4°C for further analysis. The proliferation of MCF-7 cells was assessed by MTT assay. **RESULTS** The present study demonstrated the anticancer activity of Rhizophora mucronata mangrove leaf extract against lung cancer cell line and the result is graphically represented. **CONCLUSION** The obtained result established the anticancer activity of Rhizophora mucronata mangrove leaf extract against lung cancer cell lines.

Keywords: Mangrove plant; Rhizophora mucronata; Bioactive compounds; Lung cancer cell line

1. Introduction

Ayurveda is one of the oldest health systems in India. In Sanskrit, Ayu means life and Veda means knowledge or science. Ayurvedic medicine is based on six darshanas; it includes logic of Samkhya, Nyaya vaishesika systems of natural philosophy. Ayurveda is the most comprehensive and practical medical science that revives support of the public. Ayurvedic medicine remained stagnated and even suppressed. However, it survived against the adversity of time, especially during the Mughal and British period (Patwardhan, 2010; Rajeshkumar et al., 2018). Rhizophora mucronata is a slow-growing, much-branched, evergreen tree growing up to 27 metres tall, with a bole 50 - 70 cm in diameter. The tree produces numerous stilt roots from the base. The tree is an important local source of tannins and fuel, mainly gathered from the wild but also semi-

cultivated in some areas. Traditional Use of Rhizophora mucronata is the bark of this plant is known as an astringent. It has traditionally been used in the treatment of diabetes, diarrhoea, nausea, haematuria, haemorrhages and angina ('Indian Medicinal Plants', 2007) (Kirtikara and Basu, 1918) The traditional use of several mangrove plants including R. mucronata has recently been reviewed (Website, no date; Nandhini, Rajeshkumar and Mythili, 2019). The most use of this plant is anti-cancer activity (Premanathan, Kathiresan and Chandra, 1995) ('Indian Medicinal Plants', 2007; Santhoshkumar et al., 2019) (Premanathan et al., 1999) The ethanol bark extract was found to have high activity against the Newcastle disease, vaccinia, encephalomyocarditis and Forest viruses. Also the ethanol flower extract showed good activity in human health. The bark of Rhizophora mucronata was the most promising Anticancer agent (R et al., 2020).

Rhizophora mucronata leaf extracts exhibit best antiviral and anti-cancer natural agents. The natural compounds used in both traditional and modern therapies for human health improvement with fewer side effects (Saravanan et al., 2018; Gheena and Ezhilarasan, 2019). Leaf extracts show novel drug molecules to combat the threat of human diseases (Kuppusamy et al., 2015) (Arumugam, Swamy and Sinniah, 2016; Swamy and Sinniah, 2016). The screening of plant extracts is an innovative method to find therapeutically active compounds in many plant species (Raghunandhakumar et al., 2013; Koka et al., 2018). Our team has extensive knowledge and research experience that has translated into high quality publications (Choudhari and Thenmozhi, 2016; Govindaraju, Jeevanandan and Subramanian, 2017; Ravi et al., 2017; Vikram et al., 2017; Gupta, Ariga and Deogade, 2018; Hannah et al., 2018; Kavarthapu and Thamaraiselvan, 2018; Pandian, Krishnan and Kumar, 2018; Ramamurthy and Mg, 2018; Ashok and Ganapathy, 2019; Ramesh et al., 2019; Sharma et al., 2019; Venu, Raju and Subramani, 2019; Wu et al., 2019; Samuel, Acharya and Rao, 2020). The aim of the study is to evaluate the anti cancer activity from rhizophora mucronata mangrove leaf extracts against lung cancer cell lines.

2. Materials and Methods

Sample collection and preparation

The Rhizophora mucronata mangrove leaf samples were collected from the Parangipettai, Tamilnadu. The collected sample was washed thoroughly with tap water then was dried in shadow for 48 hrs and further oven dried for 24hrs and turned into fine powder by mortar and pestle.

Preparation of Extraction

25g of dried powdered Rhizophora mucronata sample was mixed with 100ml of Ethanol and allowed to place for 24 hours at ambient temperature. Then the mixture was passing through whatman filter paper (No.4) then the filtrate was centrifuged at 3000rpm for 10min and further filtered by 0.45µm syringe micro filter. At last, the solvents are evaporated via vacuum rotary evaporator until samples obtain in powder form. Then the sample was stored in a shadowy aluminum container at 4°C for further analysis.

MTT Assay: The proliferation of MCF-7 cells was assessed by MTT assay Safadi et al., (2003). MCF-7 cells were plated in 48 well plates at a concentration of 2x10⁴ cells/well 24 hours after plating, cells were washed twice with 500µl of serum-free medium and starved by incubating the cells in serum-free medium for 3 hours at 37°C. After starvation, cells were treated with Rhizophora mucronata extract in different concentrations for 24 hours. At the end of treatment, the medium from control and Rhizophora mucronata extract treated cells were discarded and 200µl of MTT containing DMEM (0.5 mg/ml) was added to each well. The cells were then incubated for 4h at 37°C in the CO₂ incubator.

The MTT containing medium was then discarded and the cells were washed with 1x PBS. The crystals were then dissolved by adding 200µl of solubilization solution and this was mixed properly by pipetting up and down. Then the formed crystals were dissolved in dimethylsulfoxide (200µl) and incubated in dark for an hour. Then the intensity of the color developed was assayed using a Micro ELISA plate reader at 570 nm. The number of viable cells was expressed as the percentage of control cells cultured in serum-free medium. Cell viability in control medium without any treatment was represented as 100%. The cell viability is calculated using the formula: % cell viability = [A_{570 nm} of treated cells/A_{570 nm} of control cells] × 100.

3. Morphology Study

Based on MTT assay we selected the optimal doses (300µg/ml) for further studies. Analysis of cell morphology changes by a phase contrast microscope. 3×10⁴ cells were seeded in 6 well plates and treated with R. mucronata extract (500µg/ml for MCF-7 cells) for 24h. At the end of the incubation period, the medium was removed, and cells were washed once with a phosphate buffer saline (PBS pH 7.4). The plates were observed under a phase contrast microscope.

4. Statistical Analysis

All data obtained were analyzed by Student's-t-test using MS-Excel, represented as mean ± SD for triplicates. The results were computed statistically (SPSS/10 Software Package; SPSS Inc., Chicago, IL, USA) using one-way ANOVA. The level of statistical significance was set at p<0.05.

5. Results and Discussion




Figure 1: showing anticancer activity of Rhizophora mucronata mangrove leaf extract against lung cancer cell line. X-Axis represents the concentration in µl and Y-Axis represents the percentage of cell viability.

It is shown in the results that the Rhizophora mucronata tested in this investigation possess anticancer activity. Mangrove leaf extract have high phenolics and flavonoid contents and also, high antioxidant activity with a low IC₅₀. Also, there are strong positive and significant correlations between DPPH radical scavenging and phenolics and flavonoid contents. As the phenolic compounds and flavonoids are the main contributors of anticancer

activity in these mangrove leaf extract. (Fig 1)
 The contribution of new and novel products from potential bioactive plants or their extracts for disease treatment and prevention is still vast, despite the overshadowing by recent synthetic chemistry as a method of drug discoveries and drug productions (Raghunandhakumar et al., 2013)(Asokkumar et al., 2012). Moreover, plant derived drugs like vinblastine, vincristine, taxol, and camptothecin had led to greatest extent within the vicinity of antitumor upon where the drugs were reported to improvise the chemotherapy of some cancers (Yousefzadi et al., 2010). Plants contain almost unlimited capacity to generate compounds that fascinates researchers in the quest for new and novel chemotherapeutics. The persistence search for new Anticancer compounds in plant medicines and traditional foods is a realistic and promising strategy for its prevention. Numerous compounds found in plants with anticancer properties are such as alkaloids, phenylpropanoids, and terpenoids. Therefore, in this study Rhizophora mucronata mangrove leaf extract was evaluated as a new anticancer agent by using MTT assays. Plants used in folk and traditional medicines have been accepted as leads for therapeutic drug development in modern medicine. Rhizophora mucronata mangrove leaf extract was chosen for this study due to its use as a wound healing agent among the natives of Africans and as therapeutic agent in other parts of the world (Ezhilarasan, Apoorva and Vardhan, 2018; Gnanavel, Roopan and Rajeshkumar, 2019; Markov et al., 2021). Hence this study the cytotoxicity was evaluated in vitro. Studies have observed the presence of a large number of bioactive compounds in the methanolic extracts of this plant including tannins, alkaloids, steroids, saponins, terpenoids, and flavonoids which exhibit various biological activities[(Gulecha and Sivakuma, 2011)(Kumar, Jaya Kumar and Jaya Santhi, 2012)(Hussain et al., 2012)(Kumar, Rajkapoor and Perumal, 2011) These compounds are present in a number of food items and hold great potential as drug candidates due to their safety, low toxicity and wide acceptance amongst the public.(Ezhilarasan, 2018; Ezhilarasan, Sokal and Najimi, 2018; Gomathi et al., 2020)

The focus of the current research has been based on the identification of natural and synthetic compounds that can be used in the prevention or the treatment of cancer(Ezhilarasan, Apoorva and Vardhan, 2018; Duraisamy et al., 2019; Joseph and Prasanth, 2021). Methanolic extract of Rhizophora mucronata showed the presence of carbohydrates, phenol, alkaloid, amino acids, fat, flavonoids, glycosides, phenols, protein, saponins, sterols and tannins. Fourier-transform infrared spectroscopy (FTIR) confirmed the presence of primary, secondary amines, amides, alkanes, alkynes, aldehydes, saturated aliphatic, primary amines, alkyl halides, aliphatic amines, carboxylic acids, alkyl halides and alkyne(Dinesh and Ramanathan, 2017)(Ramesh et al., 2018; Dua et al., 2019; Arumugam, George and

Jayaseelan, 2021). The percentage of viability of Michigan Cancer Foundation-7 (MCF-7) cell line for control is 100%. Based on the result of present study, it can also be concluded that the effect on methanolic extract of *R. mucronata* has the anticancer activity potential in MCF-7 cell line. The limitations can be constrained by global climate change and terrestrial for its vegetation. The future scope is to create awareness and better knowledge about the anticancer properties of Rhizophora mucronata as found that high dietary intake of natural phenolics is strongly associated with longer life expectancy, reduced risk of developing some chronic diseases, various types of cancer, diabetes, obesity, improved endothelial function and reduced blood pressure

6. Conclusion

In conclusion, the obtained result established the preliminary phytochemical analysis of Rhizophora mucronata leaf. The outcome of the existing study encourages carrying out further compound's identification present in the extract. Large screening is required for the evaluation of natural products. This provides a way for the future pipeline in drug discovery. Thus, Rhizophora mucronata possesses various pharmaceutical properties in the field of herbal medicine.

7. Acknowledgement

The authors would like to thank Saveetha Dental College, SIMATS for providing a platform to conduct this research.

8. Conflict of Interest

The authors declare that there was no conflict of interest

References

- Arumugam, G., Swamy, M. and Sinniah, U. (2016) 'Plectranthus amboinicus (Lour.) Spreng: Botanical, Phytochemical, Pharmacological and Nutritional Significance', *Molecules*, p. 369. doi:10.3390/molecules21040369.
- Arumugam, P., George, R. and Jayaseelan, V.P. (2021) 'Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma', *Archives of oral biology*, 122, p. 105030. doi:10.1016/j.archoralbio.2020.105030.
- Ashok, V. and Ganapathy, D. (2019) 'A geometrical method to classify face forms', *Journal of oral biology and craniofacial research*, 9(3), pp. 232–235. doi:10.1016/j.jobcr.2019.06.001.
- Asokkumar, S. et al. (2012) 'Antiproliferative and antioxidant potential of beta-ionone against benzo(a)pyrene-induced lung carcinogenesis in Swiss albino mice', *Molecular and cellular biochemistry*, 363(1-2), pp. 335–345. doi:10.1007/s11010-011-1186-6.

- Choudhari, S. and Thenmozhi, M.S. (2016) 'Occurrence and Importance of Posterior Condylar Foramen', *Journal of advanced pharmaceutical technology & research*, 9(8), p. 1083. doi:10.5958/0974-360x.2016.00206.7.
- Dinesh, P. and Ramanathan, T. (2017) 'Anti-cancer Activities of Methanol Extract of Rhizophora mucronata against Breast Cancer MCF7 Cell Line', *Indian Internet Journal of Forensic Medicine & Toxicology*, p. 71. doi:10.5958/0974-4487.2017.00013.x.
- Dua, K. et al. (2019) 'The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress', *Drug Development Research*, pp. 714–730. doi:10.1002/ddr.21571.
- Duraisamy, R. et al. (2019) 'Compatibility of Nonoriginal Abutments With Implants', *Implant Dentistry*, pp. 289–295. doi:10.1097/id.0000000000000885.
- Ezhilarasan, D. (2018) 'Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective', *Arab Journal of Gastroenterology*, pp. 56–64. doi:10.1016/j.ajg.2018.03.002.
- Ezhilarasan, D., Apoorva, V.S. and Vardhan, N.A. (2018) 'Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of Oral Pathology & Medicine* [Preprint]. doi:10.1111/jop.12806.
- Ezhilarasan, D., Sokal, E. and Najimi, M. (2018) 'Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets', *Hepatobiliary & Pancreatic Diseases International*, pp. 192–197. doi:10.1016/j.hbpd.2018.04.003.
- Gheena, S. and Ezhilarasan, D. (2019) 'Synergic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & Experimental Toxicology*, pp. 694–702. doi:10.1177/0960327119839173.
- Gnanavel, V., Roopan, S.M. and Rajeshkumar, S. (2019) 'Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products', *Aquaculture*, pp. 1–6. doi:10.1016/j.aquaculture.2019.04.004.
- Gomathi, A.C. et al. (2020) 'Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of Tamarindus indica on MCF-7 human breast cancer cell line', *Journal of Drug Delivery Science and Technology*, p. 101376. doi:10.1016/j.jddst.2019.101376.
- Govindaraju, L., Jeevanandan, G. and Subramanian, E. (2017) 'Clinical Evaluation of Quality of Obturation and Instrumentation Time using Two Modified Rotary File Systems with Manual Instrumentation in Primary Teeth', *Journal of clinical and diagnostic research: JCDR*, 11(9), pp. ZC55–ZC58. doi:10.7860/JCDR/2017/30069.10602.
- Gulecha, V. and Sivakuma, T. (2011) 'Anticancer activity of Tephrosia purpurea and Ficus religiosa using MCF 7 cell lines', *Asian Pacific Journal of Tropical Medicine*, pp. 526–529. doi:10.1016/s1995-7645(11)60139-9.
- Gupta, P., Ariga, P. and Deogade, S.C. (2018) 'Effect of Monopoly-coating Agent on the Surface Roughness of a Tissue Conditioner Subjected to Cleansing and Disinfection: A Contact Profilometric In vitro Study', *Contemporary clinical dentistry*, 9(Suppl 1), pp. S122–S126. doi:10.4103/ccd.ccd_112_18.
- Hannah, R. et al. (2018) 'Awareness about the use, ethics and scope of dental photography among undergraduate dental students dentist behind the lens', *Journal of advanced pharmaceutical technology & research*, 11(3), p. 1012. doi:10.5958/0974-360x.2018.00189.0.
- Hussain, T. et al. (2012) 'Acute and subacute oral toxicity evaluation of Tephrosia purpurea extract in rodents', *Asian Pacific Journal of Tropical Disease*, pp. 129–132. doi:10.1016/s2222-1808(12)60030-9.
- 'Indian Medicinal Plants' (2007). doi:10.1007/978-0-387-70638-2.
- Joseph, B. and Prasanth, C.S. (2021) 'Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry?', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, pp. 118–119. doi:10.1016/j.oooo.2021.01.025.
- Kavarthapu, A. and Thamaraiselvan, M. (2018) 'Assessing the variation in course and position of inferior alveolar nerve among south Indian population: A cone beam computed tomographic study', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(4), pp. 405–409. doi:10.4103/ijdr.IJDR_418_17.
- Kirtikara, K.R. and Basu, B.D. (1918) 'Indian medicinal plants /'. doi:10.5962/bhl.title.137025.
- Koka, P. et al. (2018) 'Uncoupling Warburg effect and stemness in CD133 cancer stem cells from Saos-2 (osteosarcoma) cell line under hypoxia', *Molecular biology reports*, 45(6), pp. 1653–1662. doi:10.1007/s11033-018-4309-2.
- Kumar, D.J., Jaya Kumar, D. and Jaya Santhi, R. (2012) 'Antioxidant and cytotoxic effects of hexane extract of Morinda pubescens leaves in human liver cancer cell line', *Asian Pacific Journal of Tropical Medicine*, pp. 362–366. doi:10.1016/s1995-7645(12)60060-1.
- Kumar, R.S., Raj Kapoor, B. and Perumal, P. (2011) 'In vitro and in vivo anticancer activity of Indigofera cassioides Rottl. Ex. DC', *Asian Pacific Journal of Tropical Medicine*, pp. 379–385. doi:10.1016/s1995-7645(11)60108-9.
- Kuppusamy, P. et al. (2015) 'Evaluation of in-vitro antioxidant and antibacterial properties of Commelina nudiflora L. extracts prepared by different polar solvents', *Saudi journal of biological sciences*, 22(3), pp. 293–301. doi:10.1016/j.sjbs.2014.09.016.
- Markov, A. et al. (2021) 'Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders', *Stem Cell Research & Therapy*. doi:10.1186/s13287-021-02265-1.
- Nandhini, N.T., Rajeshkumar, S. and Mythili, S. (2019) 'The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation', *Biocatalysis and Agricultural Biotechnology*, p.

101138. doi:10.1016/j.bcab.2019.101138.
- Pandian, K.S., Krishnan, S. and Kumar, S.A. (2018) 'Angular photogrammetric analysis of the soft-tissue facial profile of Indian adults', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(2), pp. 137–143. doi:10.4103/ijdr.IJDR_496_16.
- Patwardhan, B. (2010) 'Ayurveda, evidence-base and scientific rigor', *Journal of Ayurveda and Integrative Medicine*, p. 169. doi:10.4103/0975-9476.72605.
- Premanathan, M. et al. (1999) 'In Vitro Anti-Human Immunodeficiency Virus Activity of Polysaccharide from *Rhizophora mucronata* Poir', *Bioscience, Biotechnology, and Biochemistry*, pp. 1187–1191. doi:10.1271/bbb.63.1187.
- Premanathan, M., Kathiresan, K. and Chandra, K. (1995) 'Antiviral Evaluation of Some Marine Plants Against Semliki Forest Virus', *International Journal of Pharmacognosy*, pp. 75–77. doi:10.3109/13880209509088153.
- Raghunandhakumar, S. et al. (2013) 'Thymoquinone inhibits cell proliferation through regulation of G1/S phase cell cycle transition in N-nitrosodiethylamine-induced experimental rat hepatocellular carcinoma', *Toxicology letters*, 223(1), pp. 60–72. doi:10.1016/j.toxlet.2013.08.018.
- Rajeshkumar, S. et al. (2018) 'Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells', *Enzyme and Microbial Technology*, pp. 91–95. doi:10.1016/j.enzmictec.2018.06.009.
- Ramamurthy, J. and Mg, V. (2018) 'Comparison of effect of Hiora mouthwash versus Chlorhexidine mouthwash in gingivitis patients: A clinical trial', *Asian journal of pharmaceutical and clinical research*, 11(7), p. 84. doi:10.22159/ajpcr.2018.v11i7.24783.
- Ramesh, A. et al. (2018) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of Periodontology*, pp. 1241–1248. doi:10.1002/jper.17-0445.
- Ramesh, A. et al. (2019) 'Esthetic lip repositioning: A cosmetic approach for correction of gummy smile - A case series', *Journal of Indian Society of Periodontology*, 23(3), pp. 290–294. doi:10.4103/jisp.jisp_548_18.
- Ravi, S. et al. (2017) 'Additive Effect of Plasma Rich in Growth Factors With Guided Tissue Regeneration in Treatment of Intrabony Defects in Patients With Chronic Periodontitis: A Split-Mouth Randomized Controlled Clinical Trial', *Journal of Periodontology*, pp. 839–845. doi:10.1902/jop.2017.160824.
- R, K.R. et al. (2020) ' β -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line', *Journal of Biomedical Materials Research Part A*, pp. 1899–1908. doi:10.1002/jbm.a.36953.
- Samuel, S.R., Acharya, S. and Rao, J.C. (2020) 'School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial', *Journal of public health dentistry*, 80(1), pp. 51–60. doi:10.1111/jphd.12348.
- Santhoshkumar, J. et al. (2019) 'Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*', *South African Journal of Chemical Engineering*, pp. 17–23. doi:10.1016/j.sajce.2019.04.001.
- Saravanan, M. et al. (2018) 'Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria', *Microbial Pathogenesis*, pp. 68–72. doi:10.1016/j.micpath.2018.02.008.
- Sharma, P. et al. (2019) 'Emerging trends in the novel drug delivery approaches for the treatment of lung cancer', *Chemico-biological interactions*, 309, p. 108720. doi:10.1016/j.cbi.2019.06.033.
- Swamy, M.K. and Sinniah, U.R. (2016) 'Patchouli (*Pogostemon cablin* Benth.): Botany, agrotechnology and biotechnological aspects', *Industrial Crops and Products*, pp. 161–176. doi:10.1016/j.indcrop.2016.04.032.
- Venu, H., Raju, V.D. and Subramani, L. (2019) 'Combined effect of influence of nano additives, combustion chamber geometry and injection timing in a DI diesel engine fuelled with ternary (diesel-biodiesel-ethanol) blends', *Energy*, 174, pp. 386–406. doi:10.1016/j.energy.2019.02.163.
- Vikram, N.R. et al. (2017) 'Ball Headed Mini Implant', *Journal of clinical and diagnostic research: JCDR*, 11(1), pp. ZL02–ZL03. doi:10.7860/JCDR/2017/24358.9240.
- Website (no date). Available at: Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity [Internet]. Vol. 32, South African Journal of Chemical Engineering. 2020. p. 1–4. Available from: <http://dx.doi.org/10.1016/j.sajce.2019.11.005>.
- Wu, F. et al. (2019) 'Biologically synthesized green gold nanoparticles from *Siberian ginseng* induce growth-inhibitory effect on melanoma cells (B16)', *Artificial Cells, Nanomedicine, and Biotechnology*, pp. 3297–3305. doi:10.1080/21691401.2019.1647224.
- Yousefzadi, M. et al. (2010) 'Podophyllotoxin: Current approaches to its biotechnological production and future challenges', *Engineering in Life Sciences*, pp. 281–292. doi:10.1002/elsc.201000027.